



University of Richmond  
**UR Scholarship Repository**

---

Math and Computer Science Faculty Publications

Math and Computer Science

---

1994

# Analytic Besov Spaces and Invariant Subspaces of Bergman Spaces

William T. Ross

*University of Richmond*, [wross@richmond.edu](mailto:wross@richmond.edu)

Follow this and additional works at: <http://scholarship.richmond.edu/mathcs-faculty-publications>

 Part of the [Algebra Commons](#)

---

## Recommended Citation

Ross, William T. "Analytic Besov Spaces and Invariant Subspaces of Bergman Spaces." *Indiana University Mathematics Journal* 43, no. 4 (1994): 1297-1319. doi:[10.1512/iumj.1994.43.43056](https://doi.org/10.1512/iumj.1994.43.43056).

This Article is brought to you for free and open access by the Math and Computer Science at UR Scholarship Repository. It has been accepted for inclusion in Math and Computer Science Faculty Publications by an authorized administrator of UR Scholarship Repository. For more information, please contact [scholarshiprepository@richmond.edu](mailto:scholarshiprepository@richmond.edu).

# *Analytic Besov Spaces and Invariant Subspaces of Bergman Spaces*

WILLIAM T. ROSS

ABSTRACT. In this paper, we examine the invariant subspaces (under the operator  $f \rightarrow zf$ )  $\mathcal{M}$  of the Bergman space  $L_a^p(G \setminus \mathbb{T})$  (where  $1 < p < 2$ ,  $G$  is a bounded region in  $\mathbb{C}$  containing  $\mathbb{D}$ ,  $\mathbb{T}$  is the unit circle, and  $\mathbb{D}$  is the unit disk) which contain the characteristic functions  $\chi_{\mathbb{D}}$  and  $\chi_G$ , i.e. the constant functions on the components of  $G \setminus \mathbb{T}$ . We will show that such  $\mathcal{M}$  are in one-to-one correspondence with the invariant subspaces of the analytic Besov space  $AB_q$  ( $q$  is the conjugate index to  $p$ ) and then use results of Shirokov to describe such  $\mathcal{M}$ . When  $p \geq 2$  the situation becomes more complicated and capacity considerations are needed.

**1. Introduction.** For  $1 < p < \infty$  and a bounded open set  $U \subset \mathbb{C}$ , the Bergman space  $L_a^p(U)$  is the space of analytic functions  $f$  on  $U$  for which

$$\int_U |f(z)|^p dx dy < \infty.$$

The subspaces  $\mathcal{M} \subset L_a^p(U)$  with  $z\mathcal{M} \subset \mathcal{M}$  (We will call such subspaces *invariant subspaces*.) are so fantastically complicated that they defy a reasonable characterization. In this paper, we wish to continue an investigation begun in [2], [23] of the invariant subspaces

$$\chi_G \in \mathcal{M} \subset L_a^p(G \setminus K),$$

where  $K$  is a compact subset of a bounded region  $G \subset \mathbb{C}$  and  $\text{Area}(K) = 0$ . When  $1 < p < 2$  and  $G \setminus K$  is connected,  $\mathcal{M}$  has a relatively simple characterization as  $\mathcal{M} = L_a^p(G \setminus E)$  for some closed  $E \subset K$ . When  $p \geq 2$  and  $G \setminus K$  is connected,

not all  $\mathcal{M}$  are of the form  $L_a^p(G \setminus E)$  but instead take the form

$$\mathcal{M} = \overline{\bigcup_n L_a^p(G \setminus E_n)}^{L^p},$$

where  $\{E_n\}$  is an increasing sequence of closed subsets of  $K$ .

When  $G \setminus K$  is not connected, the problem becomes much more complicated. In this paper we begin to investigate this situation in the special case when  $G$  is a region containing the closure of the open unit disk  $\mathbb{D}$ ,  $K$  is the unit circle  $\mathbb{T}$ , and the invariant subspace  $\mathcal{M}$  has the property

$$\chi_G, \chi_{\mathbb{D}} \in \mathcal{M}.$$

(i.e.  $\mathcal{M}$  contains the constants on the components of  $G \setminus \mathbb{T}$ ).

**Remark.** Throughout this paper, a ‘region’ will be an open connected subset of the plane and a ‘domain’ will be a open subset of the plane (it need not be connected).

It will turn out, via annihilators and the Cauchy transform, that such invariant subspaces  $\mathcal{M}$  will be in one-to-one correspondence with the invariant subspaces (under multiplication by  $\zeta$ ) of the analytic Besov space  $AB_q$  ( $q$  is the conjugate index to  $p$ ) of Hardy space  $H^q$  functions with

$$\int_{\mathbb{T}} \int_{\mathbb{T}} \left| \frac{f(\zeta) - f(\xi)}{\zeta - \xi} \right|^q |d\zeta| |d\xi| < \infty.$$

When  $1 < p < 2$ , then  $q > 2$  and  $AB_q$  becomes an Banach algebra of continuous functions on  $\mathbb{T}$  and the invariant subspaces are the closed ideals of  $AB_q$  which have been characterized by Shirokov [26] as

$$\mathcal{I} = \mathcal{I}(E, I) \equiv \{f \in AB_q : f|_E = 0, f/I \in H^\infty\},$$

for some closed set  $E \subset \mathbb{T}$  and inner function  $I \in H^\infty(\mathbb{D})$ . Moreover, if  $I = BS_\mu$  is the usual factorization of the inner function  $I$  into a Blaschke product  $B$ , with zeros  $\{a_k\}$ , and a singular inner function  $S_\mu$ , with positive singular measure  $\mu$ , then we set

$$\text{spec}(I) = \text{clos}\{a_k\} \cup \text{supp}(\mu).$$

With this notation, it is known (see below) that the ideal  $\mathcal{I}(E, I) \neq (0)$  if and only if the following condition is satisfied:

$$(1.1) \quad \int_{\mathbb{T}} \log \text{dist}(\zeta, E \cup \text{spec}(I)) |d\zeta| > -\infty.$$

Thus for  $1 < p < 2$ , every invariant subspace  $\chi_{\mathbb{D}}, \chi_G \in \mathcal{M} \subset L_a^p(G \setminus \mathbb{T})$  can be written as  $\mathcal{M}_{\mathcal{F}(E,I)}$  and we will show that  $\mathcal{M}_{\mathcal{F}(E,I)} = L_a^p(G \setminus \mathbb{T})$  if and only if  $\mathcal{F}(E,I) = 0$ . Our first theorem identifies  $\mathcal{M}_{\mathcal{F}(E,I)}$ .

**Theorem 1.1.** *Let  $1 < p < 2$  and  $\chi_G, \chi_{\mathbb{D}} \in \mathcal{M} \subset L_a^p(G \setminus \mathbb{T})$  be invariant. Then there is a closed set  $E \subset \mathbb{T}$  and an inner function  $I \in H^\infty(\mathbb{D})$  with*

$$\mathcal{M} = \mathcal{M}_{\mathcal{F}(E,I)} = L_a^p(G \setminus E) \vee \left\{ \frac{\chi_{G \setminus \mathbb{D}}}{\phi} L_a^p(G) : \phi \text{ inner}, \frac{I}{\phi} \in H^\infty \right\}$$

Moreover,  $\mathcal{M} \neq L_a^p(G \setminus \mathbb{T})$  if and only if condition (1.1) is satisfied.

Here we use the notation  $A \vee B$  to denote the closed linear span of  $A$  and  $B$ . Notice that for an inner function  $\phi$  and  $|z| > 1$ , we have  $\phi(z) = \phi(z^*)^*$  ( $a^* = 1/\bar{a}$ ) and hence  $1/|\phi(z)| \leq 1$  for  $|z| > 1$ . Thus  $\chi_{G \setminus \mathbb{D}}/\phi \in L_a^p(G \setminus \mathbb{T})$ . Throughout this paper when we use the term inner function, we mean a bounded analytic function on the unit disk  $\mathbb{D}$  which is unimodular a.e. on the unit circle  $\mathbb{T}$  (in contrast to a inner function defined on a general domain). We can refine Theorem 1.1 as follows: If the inner function  $I$  is the least common multiple (see definition below) of the inner functions  $I_1$  and  $I_2$  (e.g.  $I_1 = 1, I_2 = I$ , or  $I_1 = B, I_2 = S_\mu$ ), then

$$\mathcal{M}_{\mathcal{F}(E,I)} = L_a^p(G \setminus E) \vee \frac{\chi_{G \setminus \mathbb{D}}}{I_1} L_a^p(G) \vee \frac{\chi_{G \setminus \mathbb{D}}}{I_2} L_a^p(G).$$

For  $p = 2$ , the problem becomes more complicated since the analytic Besov space  $AB_2$  (often called the Dirichlet space) is no longer an algebra of continuous (or even bounded) functions and the invariant subspaces of are not completely understood. It is a result of Beurling [6] that for  $f \in AB_2$ , the radial limit

$$\lim_{r \rightarrow 1} f(r\zeta)$$

exists quasi-everywhere (q.e.), that is to say everywhere except possibly on a set of Bessel capacity (see definition below) zero. For a set  $E \subset \mathbb{T}$ , define  $AB_{2,E}$  to be the set of  $f \in AB_2$  with radial limit zero q.e. on  $E$ . One shows [7] that  $AB_{2,E}$  is a closed invariant subspace of  $AB_2$  and it is an open question as to whether or not all invariant subspaces of  $AB_2$  are of the form

$$\mathcal{F}_{E,I} \equiv IH^2 \cap AB_{2,E}$$

for some inner function  $I$  and set  $E \subset \mathbb{T}$ . We will show, as in the  $q > 2$  case, that there is a one-to-one correspondence between the invariant subspaces of  $AB_2$  and the invariant subspaces

$$\chi_{\mathbb{D}}, \chi_G \in \mathcal{M} \subset L_a^2(G \setminus \mathbb{T}).$$

We will then identify  $\mathcal{M}_{\mathcal{F}_{E,I}}$ . Thus a complete description of the invariant subspaces of  $AB_2$  as  $\mathcal{F}_{E,I}$  will yield a complete description of the invariant subspaces  $\mathcal{M}$ . These results generalize to  $p > 2$ .

Before proceeding, we mention that the condition requiring both  $\chi_G$  and  $\chi_{\mathbb{D}}$  belong to  $\mathcal{M}$  is not a superfluous one. Without it, the problems becomes nearly impossible to solve as can be seen by the following example: By [4], Corollary 6.9 and Proposition 5.4, given any  $n \in \mathbb{N} \cup \{\infty\}$  there is an invariant subspace  $\mathcal{N}_n$  of  $L_a^2(\mathbb{D})$  with  $\dim(\mathcal{N}_n/z\mathcal{N}_n) = n$ . Consider the invariant subspace

$$(1.2) \quad \mathcal{M}_n = \chi_{\mathbb{D}}\mathcal{N}_n + L_a^2(G).$$

One shows that  $\mathcal{M}_n$  is closed in  $L_a^2(G \setminus \mathbb{T})$  and that

$$\dim(\mathcal{M}_n/z\mathcal{M}_n) = \dim(\mathcal{N}_n/z\mathcal{N}_n) + \dim(L_a^2(G)/zL_a^2(G)) = n + 1,$$

making  $\mathcal{M}_n$  difficult to understand. By adding the condition  $\chi_{\mathbb{D}} \in \mathcal{M}$ , we avoid such pathologies as  $\mathcal{M}_n$ .

## 2. Preliminaries.

**2.1. Sobolev and Besov spaces.** Throughout this paper,  $G$  will be a Jordan region in the complex plane  $\mathbb{C}$  (We make this restriction on  $G$  to avoid needless technicalities),  $\mathbb{D} = \{z : |z| < 1\}$ , and  $\mathbb{T} = \{z : |z| = 1\}$ . For the moment, we let  $1 < p < 2$  and  $q$  be the conjugate index to  $p$  (so  $q > 2$ ). The dual of  $L^p(G) = L^p(G, dA)$ , where  $dA$  is area measure, will be identified with  $L^q(G)$  via the bilinear pairing

$$(2.1) \quad \langle f, g \rangle = \int_G fg \, dA.$$

Define the Sobolev space  $W_1^{q,0}(G)$  as the closure of  $C_0^\infty(G)$  (infinitely differentiable functions with compact support in  $G$ ) in the norm

$$\|f\|_q = \left( \int |\nabla f|^q \, dA \right)^{1/q}.$$

Since  $q > 2$ , the Sobolev imbedding theorem yields that  $W_1^{q,0}(G)$  is a Banach algebra of continuous functions [1], p. 115. Here we mean that every function has a continuous representative. The following describes  $W_1^{q,0}(G \setminus E)$ , where  $E$  is closed, in terms of zero sets. We refer the reader to [3] for a proof.

**Proposition 2.1.** *For  $q > 2$ ,  $W_1^{q,0}(G \setminus E) = \{f \in W_1^{q,0}(G) : f|_E = 0\}$ .*

This next result of Havin [12] will be used later and relates the Bergman and Sobolev spaces.

**Lemma 2.2. (Havin)** *Let  $U$  be a bounded open set and  $1 < p < \infty$ . Then  $f \in L^q(U)$  satisfies*

$$\int_U u f \, dA = 0 \quad \forall u \in L^p_a(U)$$

*if and only if there is an  $F \in W^{q,0}_1(U)$  with  $\bar{\partial}F = f$ .*

Define the *Besov space*  $B_q$  as the space of functions  $f$  on  $\mathbb{T}$  with finite norm

$$\|f\|_{B_q} = \|f\|_{L^q(\mathbb{T}, |d\zeta|)} + \left( \int_{\mathbb{T}} \int_{\mathbb{T}} \left| \frac{f(\zeta) - f(\xi)}{\zeta - \xi} \right|^q |d\zeta| |d\xi| \right)^{1/q},$$

and note (since  $q > 2$ ) that  $B_q$  can be continuously embedded into  $\text{Lip}_{1-2/q}(\mathbb{T})$  and hence  $B_q$  is a Banach algebra of continuous functions on  $\mathbb{T}$  [5]. Define the *analytic Besov space*

$$AB_q = B_q \cap H^q,$$

where  $H^q$  is the usual Hardy space.

**Remark.** It is known [17] (see [28], Chapter 5, Section 5) that the analytic extension of  $f \in AB_q$ , given by the Poisson kernel, belongs to the  $L^q$ -Dirichlet space  $D_q$  of analytic functions on  $\mathbb{D}$  with

$$\|f\|_{D_q} = \|f\|_{L^q(\mathbb{T}, |d\zeta|)} + \left( \int_{\mathbb{D}} |f'(z)|^q \, dA(z) \right)^{1/q} < \infty$$

and moreover, the boundary values of  $f \in D_q$  on  $\mathbb{T}$  belongs to  $AB_q$  with the  $L^q$ -Dirichlet norm equivalent to the Besov norm. Thus we may identify  $f(\zeta) \in AB_q$  with its analytic extension  $f(z) \in D_q$ .

The spaces  $W^{q,0}_1(G)$  and  $B_q$  are related through restriction and extension. By standard trace theory [16], p. 182, [18], the trace operator

$$T : W^{q,0}_1(G) \rightarrow B_q, \quad (Th)(\zeta) = h(\zeta)$$

is a well defined, continuous, surjective linear operator with, by Proposition 1.,  $\ker(T) = W^{q,0}_1(G \setminus \mathbb{T})$ . (Note that  $h(\zeta)$  is well defined since  $q > 2$  and so  $W^{q,0}_1(G)$  is a space of continuous functions.) Thus  $T$  will induce the continuous invertible operator

$$(2.2) \quad \tilde{T} : W^{q,0}_1(G)/W^{q,0}_1(G \setminus \mathbb{T}) \rightarrow B_q, \quad \tilde{T}[h](\zeta) = h(\zeta),$$

where  $[h]$  is a coset of  $W_1^{q,0}(G)/W_1^{q,0}(G\backslash\mathbb{T})$ . By the above remark

$$(2.3) \quad \tilde{T}^{-1}(AB_q) = \{f \in W_1^{q,0}(G) : f|_{\mathbb{T}} \in AB_q\} / W_1^{q,0}(G\backslash\mathbb{T}).$$

**2.2. Ideals of the Besov space.** Since  $AB_q$  ( $q > 2$ ) is a Banach algebra and analytic polynomials are dense, the  $\zeta$ -invariant subspaces are precisely the ideals of  $AB_q$  and have been characterized by Shirokov [26] as follows:

**Theorem 2.3.** *If  $\mathcal{F}$  is closed ideal of  $AB_q$ , there is a closed set  $E \subset \mathbb{T}$  and an inner function  $I$  with*

$$\mathcal{F} = \mathcal{F}(E, I) \equiv \{f \in AB_q : f|_E = 0, f/I \in H^\infty\}.$$

**Remark.**

- (i) The set  $E$  is the common zeros of  $\mathcal{F}$  and the inner function  $I$  is the greatest common divisor of the inner parts of the functions in  $\mathcal{F}$  [15], p. 85.
- (ii) If  $f \in AB_q$  with  $f/I \in H^\infty$ , then  $f/I \in AB_q$  with  $\|f/I\|_{B_q} \leq C\|f\|_{B_q}$ . In fact, division by the inner factor is a continuous operator on other spaces of ‘smooth’ functions on the disk [13].
- (iii) From basic Hardy space theory [15], Chapter 5, an inner function  $I$  can be factored as  $I = BS_\mu$ , where  $B$  is a Blaschke product with zeros  $\{a_k\} \subset \mathbb{D}$  (repeated according to multiplicity) and  $S_\mu$  is a singular inner function with positive singular measure  $\mu$ . Moreover [15], p. 68–69,  $S_\mu$  cannot be continuously extended from  $\mathbb{D}$  to any point in the support of  $\mu$ . Hence, if  $f \in AB_q$  with  $f/I \in AB_q$ , then  $f$  must vanish on the support of  $\mu$  as well as the closure of  $\{a_k\}$ , i.e. on  $\text{spec}(I)$ .

For a general closed set  $E \subset \mathbb{T}$  and inner function  $I$ , the ideal  $\mathcal{F}(E, I)$  might be zero. To understand when this happens, we make the following definition: A closed set  $E \subset \mathbb{T}$  is called a *Carleson set* if

$$\int_{\mathbb{T}} \log \text{dist}(\zeta, E) |d\zeta| > -\infty.$$

It is clear from the above condition that  $E$  has Lebesgue measure zero and it is known [7] that a  $E$  is a Carleson set if and only if  $E$  has Lebesgue measure zero and

$$\sum_n |I_n| \log |I_n| > -\infty,$$

where  $\{I_n\}$  are the complimentary arcs of  $E$ .

**Proposition 2.4.** *The ideal  $\mathcal{F}(E, I)$  is non-zero if and only if condition (1.1) is satisfied.*

*Proof.* The proof is essentially known (and true for other ideals of analytic functions [29] [30]) but not explicitly stated for the Besov space, so we outline it here. If condition (1.1) is satisfied, then we can find a non-zero  $\phi \in A^\infty$  ( $f^{(n)}$  analytic on  $\mathbb{D}$  and continuous on  $\bar{\mathbb{D}} \ \forall n \in \mathbb{N} \cup \{0\}$ ) with  $\phi^{-1}(0) \cap \mathbb{T} = E$  and  $\phi/B \in A^\infty$  [29] [30], Theorem 4.1. Notice that condition (1.1) implies that the support of  $\mu$  is a Carleson set and we thus can apply [30], Corollary 4.8, to obtain a non-zero  $\psi \in A^\infty$  with  $\psi/S_\mu \in A^\infty$ . Thus  $0 \neq \phi\psi \in \mathcal{F}(E, BS_\mu)$ .

For the converse, it is known that if  $f \in AB_q$  ( $q > 2$ ), then  $f$  satisfies the Lipschitz condition [1], p. 97–98,

$$(2.4) \quad |f(z) - f(w)| \leq C|z - w|^{1-2/q} \quad z, w \in \bar{\mathbb{D}}.$$

In particular, if  $f$  is a non-zero element of  $\mathcal{F}(E, BS_\mu)$ , then by Jensen's inequality [15], p. 51–52,

$$\int_{\mathbb{T}} \log |f(\zeta)| |d\zeta| > -\infty$$

and by (2.4), along with the fact that  $f$  vanishes on  $E \cup \text{spec}(I)$ ,

$$\log |f(\zeta)| \leq (1 - 2/q) \log \text{dist}(\zeta, E \cup \text{spec}(I)) + C.$$

Hence condition (1.1) must be satisfied. □

**3. The correspondence.** We now relate our invariant subspaces of the Bergman space with the invariant subspaces of the Besov space. If  $\chi_G \in \mathcal{M} \subset L_a^p(G \setminus \mathbb{T})$  is invariant, then  $\mathcal{M}$  contains the polynomials and hence (since  $G$  is a Jordan region and polynomials are dense in  $L_a^p(G)$ )

$$L_a^p(G) \subset \mathcal{M} \subset L_a^p(G \setminus \mathbb{T}).$$

Thus,

$$L_a^p(G \setminus \mathbb{T})^\perp \subset \mathcal{M}^\perp \subset L_a^p(G)^\perp$$

with, by our bilinear pairing (2.1),  $z\mathcal{M}^\perp \subset \mathcal{M}^\perp$ . Here for a set  $X \subset L^p(G)$  we let

$$X^\perp = \{g \in L^q(G) : \langle f, g \rangle = 0 \ \forall f \in X\}.$$

Thus, there is a one-to-one correspondence between the invariant subspaces

$$\chi_G \in \mathcal{M} \subset L_a^p(G \setminus \mathbb{T})$$



and the  $R$ -invariant subspaces of the quotient space

$$L_a^p(G)^\perp / L_a^p(G \setminus \mathbb{T})^\perp, \quad R[g] = [zg].$$

Our first result (which is also found in [2]) says that  $R$  is similar to  $M_\zeta$  (multiplication by  $\zeta$ ) on the Besov space  $B_q$ . We will include a proof here so we can refer to parts of it later.

**Theorem 3.1.** *The linear transformation*

$$J : L_a^p(G)^\perp / L_a^p(G \setminus \mathbb{T})^\perp \rightarrow B_q$$

defined by

$$J[g](\zeta) = -\frac{1}{\pi} \int_G \frac{g(z)}{z - \zeta} dA(z)$$

is a continuous invertible operator with  $JR = M_\zeta J$ . Thus there is a one-to-one correspondence between the invariant subspaces  $\chi_G \in \mathcal{M} \subset L_a^p(G \setminus \mathbb{T})$  and the lattice of  $\zeta$ -invariant subspaces of  $B_q$ .

*Proof.* For any bounded open set  $U$ , we can apply Havin's Lemma (Lemma 2.2) and the Calderon-Zygmund theory [3], p. 266, to get that the operator

$$\bar{\partial} : W_1^{q,0}(U) \rightarrow L_a^p(U)^\perp$$

is continuous and invertible with inverse given by the Cauchy transform

$$(3.1) \quad (\bar{\partial}^{-1}g)(w) = (Cg)(w) = -\frac{1}{\pi} \int_U \frac{g(z)}{z - w} dA(z).$$

If  $R_z$  is multiplication by  $z$  on  $L_a^p(U)^\perp$  and  $M_z$  is multiplication by  $z$  on  $W_1^{q,0}(U)$  (both well defined and continuous) then, noticing that  $\bar{\partial}(zf) = z\bar{\partial}f$  for all  $f \in W_1^{q,0}(U)$ , we have

$$(3.2) \quad R_z \bar{\partial} = \bar{\partial} M_z.$$

The Cauchy transform  $C = \bar{\partial}^{-1}$  will induce the continuous invertible operator

$$\tilde{C} : L_a^p(G)^\perp / L_a^p(G \setminus \mathbb{T})^\perp \rightarrow W_1^{q,0}(G) / W_1^{q,0}(G \setminus \mathbb{T}).$$

Notice that  $R_z$  and  $M_z$  will induce the multiplication operators  $R$  and  $M$  on the cosets of  $L_a^p(G)^\perp / L_a^p(G \setminus \mathbb{T})^\perp$  and  $W_1^{q,0}(G) / W_1^{q,0}(G \setminus \mathbb{T})$  respectively with, by (3.2),

$$\tilde{C}R = M\tilde{C}.$$

Thus if we define

$$J : L_a^p(G)^\perp / L_a^p(G \setminus \mathbb{T})^\perp \rightarrow B_q$$

by  $J = \tilde{T} \circ \tilde{C}$  (recall the definition of  $\tilde{T}$  in (2.2)), we obtain

$$J[g](\zeta) = -\frac{1}{\pi} \int_G \frac{g(z)}{z - \zeta} dA(z)$$

and  $JR = M_\zeta J$ , where  $M_\zeta$  is multiplication by  $\zeta$  on  $B_q$ .  $\square$

**Corollary 3.2.** *If  $\chi_{\mathbb{D}}, \chi_G \in \mathcal{M} \subset L_a^p(G \setminus \mathbb{T})$  is invariant and  $g \in \mathcal{M}^\perp$ , then*

$$J[g](\zeta) = -\frac{1}{\pi} \int_{G \setminus \mathbb{D}} \frac{g(z)}{z - \zeta} dA(z).$$

*The function  $J[g] \in AB_q$  and hence there is a one-to-one correspondence between the invariant subspaces  $\chi_{\mathbb{D}}, \chi_G \in \mathcal{M}$  and the ideals of the analytic Besov space  $AB_q$ .*

*Proof.* If  $\chi_G, \chi_{\mathbb{D}} \in \mathcal{M} \subset L_a^p(G \setminus \mathbb{T})$ , then by the invariance of  $\mathcal{M}$ , we get

$$L_a^p(G) \bigvee \chi_{\mathbb{D}} L_a^p(\mathbb{D}) \subset \mathcal{M} \subset L_a^p(G \setminus \mathbb{T}).$$

Taking annihilators one obtains

$$L_a^p(G \setminus \mathbb{T})^\perp \subset \mathcal{M}^\perp \subset L_a^p(G)^\perp \cap (\chi_{\mathbb{D}} L_a^p(\mathbb{D}))^\perp.$$

Thus if  $g \in \mathcal{M}^\perp$ , then  $Cg \in W_1^{q,0}(G)$  and since  $\mathcal{M}^\perp \subset (\chi_{\mathbb{D}} L_a^p(\mathbb{D}))^\perp$ , we have

$$0 = \langle g, \chi_{\mathbb{D}}(z - \lambda)^{-1} \rangle = \int_{\mathbb{D}} \frac{g}{z - \lambda} dA$$

for all  $|\lambda| \geq 1$  (Note that  $(z - \lambda)^{-1} \in L_a^p(\mathbb{D})$  for  $1 < p < 2$ ). Thus

$$J[g](\zeta) = (TCg)(\zeta) = -\frac{1}{\pi} \int_{G \setminus \mathbb{D}} \frac{g}{z - \zeta} dA.$$

The above function belongs to  $B_q$  and is analytic on  $\mathbb{D}$ , hence  $J[g] \in AB_q$ .  $\square$

**Notation.** If  $\mathcal{F}$  is an ideal of  $AB_q$  we let  $\mathcal{M}_{\mathcal{F}}$  be the unique invariant subspace of  $L_a^p(G \setminus \mathbb{T})$  which contains  $\chi_G$  and  $\chi_{\mathbb{D}}$  and corresponds to  $\mathcal{F}$  via  $J$ . One checks that

$$(3.3) \quad \mathcal{M}_{\mathcal{F}} = (\bar{\partial}\{f \in W_1^{q,0}(G) : f|_{\mathbb{T}} \in \mathcal{F}\})^{\perp}.$$

**4. Invariant subspaces of Bergman spaces.** Before proceeding to our main results, we first make a comment about inner functions. If  $I$  is an inner function and  $I = BS_{\mu}$ , then  $I$  is analytic for all points in the complex plane with the exception of the support of  $\mu$ ,  $\{1/\bar{a}_k\}$ , and the accumulation points of  $\{a_k\}$ . Moreover if  $w^* = 1/\bar{w}$ , then for  $|z| > 1$ ,  $I(z) = I(z^*)^*$ . Thus  $|I(z)|^{-1} \leq 1$  for all  $|z| > 1$  and so  $\chi_{G \setminus \mathbb{D}}/I \in L_a^p(G \setminus \mathbb{T})$ . We also define  $V(I)$  to be the set of inner functions  $\phi$  which divide  $I$ , that is  $I/\phi \in H^{\infty}$ .

**Theorem 4.1.** *Let  $1 < p < 2$  and  $\chi_G, \chi_{\mathbb{D}} \in \mathcal{M} \subset L_a^p(G \setminus \mathbb{T})$  be invariant. Then there is a closed set  $E \subset \mathbb{T}$  and an inner function  $I$  with*

$$\mathcal{M} = \mathcal{M}_{\mathcal{F}(E,I)} = L_a^p(G \setminus E) \bigvee \left\{ \frac{\chi_{G \setminus \mathbb{D}}}{\phi} L_a^p(G) : \phi \in V(I) \right\}$$

Moreover  $\mathcal{M} \neq L_a^p(G \setminus \mathbb{T})$  if and only if condition (1.1) is satisfied.

*Proof.* By (3.3), the unique subspace  $\mathcal{M}_{\mathcal{F}(E,I)}$  corresponding to  $\mathcal{F}(E, I)$  is

$$\mathcal{M}_{\mathcal{F}(E,I)} = (\bar{\partial}\{f \in W_1^{q,0}(G) : f|_{\mathbb{T}} \in \mathcal{F}(E, I)\})^{\perp}.$$

To finish, it suffices to show

$$\begin{aligned} & (\bar{\partial}\{f \in W_1^{q,0}(G) : f|_{\mathbb{T}} \in \mathcal{F}(E, I)\})^{\perp} \\ &= L_a^p(G \setminus E) \bigvee \left\{ \frac{\chi_{G \setminus \mathbb{D}}}{\phi} L_a^p(G) : \phi \in V(I) \right\}. \end{aligned}$$

Let  $h \in L_a^p(G \setminus E)$  and  $f \in W_1^{q,0}(G)$  with  $f|_{\mathbb{T}} \in \mathcal{F}(E, I)$ . Then by Proposition 2.1,  $f \in W_1^{q,0}(G \setminus E)$  and so by Havin's Lemma, Lemma 2.2,

$$\langle \bar{\partial}f, h \rangle = 0.$$

Thus by (3.3),  $L_a^p(G \setminus E) \subset \mathcal{M}_{\mathcal{F}(E,I)}$ .

Let  $f \in W_1^{q,0}(G)$  with  $f|_{\mathbb{T}} \in \mathcal{F}(E, I)$ . Then for  $\phi \in V(I)$ ,

$$(4.1) \quad \int_G \bar{\partial}f \frac{\chi_{G \setminus \mathbb{D}}}{\phi} dA = \lim_{\varepsilon \rightarrow 0} \int_{G \setminus \{|z| < 1 + \varepsilon\}} \frac{\bar{\partial}f}{\phi} dA =$$

$$= \lim_{\varepsilon \rightarrow 0} \int_{G \setminus \{|z| < 1+\varepsilon\}} \bar{\partial} \left( \frac{f}{\phi} \right) dA.$$

By Green's theorem [31], p. 54, and the fact that  $f = 0$  on the boundary of  $G$ , this becomes

$$- \lim_{\varepsilon \rightarrow 0} \frac{1}{2i} \int_{|z|=1+\varepsilon} \frac{f}{\phi} dz.$$

By the Lebesgue dominated convergence theorem and the fact that  $f|_{\mathbb{T}}/\phi \in H^\infty$  we obtain

$$- \frac{1}{2i} \int_{\mathbb{T}} \frac{f}{\phi} dz = 0.$$

By (3.3) we get

$$\frac{\chi_{G \setminus \mathbb{D}}}{\phi} \in \mathcal{M}_{\mathcal{F}(E, I)}.$$

Using the invariance of  $\mathcal{M}_{\mathcal{F}(E, I)}$  and the density of polynomials in  $L_a^p(G)$ , we have

$$(4.2) \quad L_a^p(G \setminus E) \vee \left\{ \frac{\chi_{G \setminus \mathbb{D}}}{\phi} L_a^p(G) : \phi \in V(I) \right\} \subset \mathcal{M}_{\mathcal{F}(E, I)}.$$

Taking annihilators and then  $C = \bar{\partial}^{-1}$  of (4.2) will yield

$$(4.3) \quad C \mathcal{M}_{\mathcal{F}(E, I)}^\perp \subset C \left( L_a^p(G \setminus E)^\perp \cap \left\{ \left( \frac{\chi_{G \setminus \mathbb{D}}}{\phi} L_a^p(G) \right)^\perp : \phi \in V(I) \right\} \right).$$

To prove equality in (4.3) and thus finish the proof, we let  $g$  belong to the right hand side of (4.3) and show that  $g \in C \mathcal{M}_{\mathcal{F}(E, I)}^\perp$  by showing  $g|_{\mathbb{T}} \in \mathcal{F}(E, I)$ , see (3.3). To do this, notice from (4.3) that  $g = C(\bar{\partial}g)$  with

$$(4.4) \quad 0 = -\frac{1}{\pi} \int_G \frac{\bar{\partial}g}{z - \lambda} dA = g(\lambda) \quad \forall \lambda \in E$$

$$(4.5) \quad \int_{G \setminus \mathbb{D}} \frac{\bar{\partial}g}{I} z^n dA = 0 \quad \forall n \in \mathbb{N} \cup \{0\}.$$

By (4.5),

$$0 = \int_{G \setminus \mathbb{D}} \frac{\bar{\partial}g}{I} z^n dA =$$

$$\begin{aligned}
&= \lim_{\varepsilon \rightarrow 0} \int_{G \setminus \{|z| < 1+\varepsilon\}} \frac{\bar{\partial} g}{I} z^n dA \\
&= \lim_{\varepsilon \rightarrow 0} \int_{G \setminus \{|z| < 1+\varepsilon\}} \bar{\partial} \left( \frac{g}{I} z^n \right) dA \quad \forall n \in \mathbb{N} \cup \{0\},
\end{aligned}$$

which, by Green's Theorem [31], p. 54, becomes

$$0 = - \lim_{\varepsilon \rightarrow 0} \frac{1}{2i} \int_{|z|=1+\varepsilon} \frac{g}{I} z^n dz \quad \forall n \in \mathbb{N} \cup \{0\}.$$

By the Lebesgue dominated convergence theorem we obtain

$$0 = - \frac{1}{2i} \int_{\mathbb{T}} \frac{g}{I} z^n dz \quad \forall n \in \mathbb{N} \cup \{0\},$$

which, by the F. and M. Riesz theorem [15], p. 47, yields  $g|_{\mathbb{T}}/I \in H^\infty$ . Thus  $g|_{\mathbb{T}}/I$  belongs to  $\mathcal{F}(E, I)$  and we are done.

Finally, notice that from (3.3) and Proposition 2.1 that  $\mathcal{F}(E, I) = 0$  if and only if  $\mathcal{M}_{\mathcal{F}(E, I)} = L_a^p(G \setminus \mathbb{T})$ . So from Proposition 2.4,  $\mathcal{M}_{\mathcal{F}(E, I)} \neq L_a^p(G \setminus \mathbb{T})$  if and only if condition (1.1) is satisfied.  $\square$

We say that an inner function  $\phi$  is a *multiple* of an inner function  $\psi$  if  $\phi/\psi \in H^\infty$ . We say an inner function  $I$  is the *least common multiple* of the inner functions  $I_1$  and  $I_2$  if  $I$  is a multiple of  $I_1$  and  $I_2$  and if the inner function  $\phi$  is a multiple of  $I_1$  and  $I_2$ , then  $\phi$  is a multiple of  $I$ .

**Corollary 4.2.** *If the inner function  $I$  is the least common multiple of the inner functions  $I_1$  and  $I_2$ , then*

$$\mathcal{M}_{\mathcal{F}(E, I)} = L_a^p(G \setminus E) \bigvee \frac{\chi_{G \setminus \mathbb{D}}}{I_1} L_a^p(G) \bigvee \frac{\chi_{G \setminus \mathbb{D}}}{I_2} L_a^p(G).$$

*Proof.* As in the proof above, we need to show

$$\begin{aligned}
&(\bar{\partial}\{f \in W_1^{q,0}(G) : f|_{\mathbb{T}} \in \mathcal{F}(E, I)\})^\perp \\
&= L_a^p(G \setminus E) \bigvee \frac{\chi_{G \setminus \mathbb{D}}}{I_1} L_a^p(G) \bigvee \frac{\chi_{G \setminus \mathbb{D}}}{I_2} L_a^p(G).
\end{aligned}$$

Using the same proof as above, one shows

$$(4.6) \quad L_a^p(G \setminus E) \subset (\bar{\partial}\{f \in W_1^{q,0}(G) : f|_{\mathbb{T}} \in \mathcal{F}(E, I)\})^\perp = \mathcal{M}_{\mathcal{F}(E, I)}.$$

Letting  $f \in W_1^{q,0}(G)$  with  $f|_{\mathbb{T}} \in \mathcal{F}(E, I)$ , we see that  $f|_{\mathbb{T}}/I_1$  and  $f|_{\mathbb{T}}/I_2$  belong to  $H^\infty$ , and thus by (4.1)

$$\frac{\chi_{G \setminus \mathbb{D}}}{I_1} \quad \text{and} \quad \frac{\chi_{G \setminus \mathbb{D}}}{I_2}$$

belong to  $\mathcal{M}_{\mathcal{F}(E, I)}$ . Now apply (4.6), the invariance of  $\mathcal{M}_{\mathcal{F}(E, I)}$ , the density of polynomials in  $L_a^p(G)$ , to show

$$(4.7) \quad \mathcal{M}_{\mathcal{F}(E, I)} \supset L_a^p(G \setminus E) \bigvee \frac{\chi_{G \setminus \mathbb{D}}}{I_1} L_a^p(G) \bigvee \frac{\chi_{G \setminus \mathbb{D}}}{I_2} L_a^p(G).$$

Taking annihilators and then  $C = \bar{\partial}^{-1}$  of (4.7) will yield

$$(4.8) \quad C\mathcal{M}_{\mathcal{F}(E, I)}^\perp \subset W_1^{q,0}(G \setminus E) \cap C \left( \left( \frac{\chi_{G \setminus \mathbb{D}}}{I_1} L_a^p(G) \right)^\perp \cap \left( \frac{\chi_{G \setminus \mathbb{D}}}{I_2} L_a^p(G) \right)^\perp \right).$$

To prove equality in (4.8), and finish the proof, we let  $g$  belong to the right hand side of (4.8) and show  $g \in C\mathcal{M}_{\mathcal{F}(E, I)}^\perp$  by showing  $g|_{\mathbb{T}} \in \mathcal{F}(E, I)$ . From (4.8),  $g = C(\bar{\partial}g)$  with

$$-\frac{1}{\pi} \int_G \frac{\bar{\partial}g}{z - \lambda} dA = g(\lambda) = 0 \quad \forall \lambda \in E,$$

$$\int_{G \setminus \mathbb{D}} \frac{\bar{\partial}g}{I_1} z^n dA = 0 \quad \forall n \in \mathbb{N} \cup \{0\},$$

$$\int_{G \setminus \mathbb{D}} \frac{\bar{\partial}g}{I_2} z^n dA = 0 \quad \forall n \in \mathbb{N} \cup \{0\}.$$

Proceed as in the proof above (using the F. and M. Riesz Theorem) to show that  $g|_{\mathbb{T}}/I_1$  and  $g|_{\mathbb{T}}/I_2$  belong to  $H^\infty$  (Just replace the inner function  $I$  in (4.5) with  $I_1$  and  $I_2$  respectively.). Letting  $I_g$  be the inner part of  $g|_{\mathbb{T}}$ , we see that  $I_g$  is a multiple of both  $I_1$  and  $I_2$ . Since  $I$  is the least common multiple of  $I_1$  and  $I_2$ , then  $I_g$  must be a multiple of  $I$ , making  $g/I \in H^\infty$ . Thus  $g|_{\mathbb{T}} \in \mathcal{F}(E, I)$  and we are done.  $\square$

**5. The classical Dirichlet space.** For  $q > 2$ , every non-zero subspace of  $AB_q$  is an ideal of the form

$$IH^\infty \cap \{f \in AB_q : f|_E = 0\}$$

for some inner function  $I$  and Carleson set  $E$ . For  $q = 2$ ,  $AB_2$  is a well studied space of analytic functions which is not an algebra and whose invariant subspaces are not completely understood.  $AB_2$  is better known as the Dirichlet space  $D_2$  and are the radial limit functions of analytic  $f(z)$  on  $\mathbb{D}$  with finite Dirichlet integral

$$\int_{\mathbb{D}} |f'(z)|^2 dA(z).$$

If  $f \in AB_2$ , a theorem of Beurling [6] says that the radial limit

$$\lim_{r \rightarrow 1} f(r\zeta)$$

exists everywhere except possibly for a set of Bessel capacity  $C_2$  (see below) zero. For  $f \in AB_2$ , let

$$Z(f) = \{\zeta \in \mathbb{T} : \lim_{r \rightarrow 1} f(r\zeta) = 0\}$$

and for a set  $E \subset \mathbb{T}$ , let

$$AB_{2,E} = \{f \in AB_2 : C_2(E \setminus Z(f)) = 0\}.$$

$AB_{2,E}$  is a closed subspace of  $AB_2$  [7] as is

$$\mathcal{F}_{E,I} \equiv IH^2 \cap AB_{2,E},$$

for an inner function  $I$  and a set  $E \subset \mathbb{T}$ , and it is a conjecture that every invariant subspace of the Dirichlet space has this form.

We will show, in a similar way to the  $1 < p < 2$  case, that the invariant subspaces of  $AB_2$  are in one-to-one correspondence with the invariant subspaces

$$\chi_{\mathbb{D}}, \chi_G \in \mathcal{M} \subset L_a^2(G \setminus \mathbb{T})$$

and then identify  $\mathcal{M}_{\mathcal{F}_{E,I}}$ . To proceed, we must first take care of some technical matters.

**5.1. Capacity.** Following [3], we define the  $C_2$ -capacity of a compact set  $F$  by

$$C_2(F) = \inf \int |\nabla \phi|^2 dA,$$

where the inf is taken over all real-valued functions  $\phi \in C_0^\infty$  with  $\phi = 1$  on  $F$ . We extend this definition to arbitrary sets  $E$  by

$$C_2(E) = \sup\{C_2(F) : F \subset E, \quad F \text{ compact}\}$$

and define the *exterior capacity*  $C_2^*(E)$  of an arbitrary set  $E$  by

$$C_2^*(E) = \inf\{C_2(G) : G \supset E, \quad G \text{ open}\}.$$

A set  $E$  is said to be *capacitable* if  $C_2(E) = C_2^*(E)$ .

**Remark.** The Sobolev space  $W_1^2$  is equal to  $L_1^2$ , the space of Bessel potentials, and thus the Bessel capacity is equivalent to  $C_2^*$  [14]. We bring this to the readers attention since the literature often uses both definitions of capacity.

One notes [3] that  $C_2^*$  is a monotone, subadditive set function and that the Borel sets are capacitable. We say a set  $E$  is *quasi-closed* of given  $\varepsilon > 0$ , there is an open set  $W$  with  $C_2(W) < \varepsilon$  and  $E \setminus W$  is closed. One argues, using the fact that Borel sets are capacitable, that a quasi-closed set is capacitable, as is the difference of any two quasi-closed sets. As mentioned in the introduction, we say a property holds *quasi-everywhere* if the set for which it fails has exterior capacity zero.

Since functions in  $W_1^{2,0}(G)$  are not always continuous (or even bounded), we introduce a suitable substitution for continuity. A complex-valued function  $f$  is *quasi-continuous* if for every  $\varepsilon > 0$  there is an open set  $W$  with  $C_2(W) < \varepsilon$  and  $f|_{\mathbb{C} \setminus W}$  continuous. One can show [3], Lemma 1, Theorem 2, that every  $f \in W_1^{2,0}(G)$  has a quasi-continuous representative and in fact, one can find a formula for the quasi-continuous representative of a Sobolev function. For  $f \in W_1^{2,0}(G)$  we define

$$(5.1) \quad f^*(w) = \lim_{r \rightarrow 0} \frac{1}{\pi r^2} \int_{|z-w| < r} f(z) dA(z)$$

whenever this limit exists and notice by the Lebesgue differentiation theorem,  $f = f^*$  a.e. By [11],  $f^*(w)$  is defined quasi-everywhere and moreover  $f^*$  is quasi-continuous. This next result of Bagby [3], Theorem 4, describes  $W_1^{2,0}(G \setminus E)$ ,  $E$  closed, in terms of zero sets.

**Proposition 5.1.**  $W_1^{2,0}(G \setminus E) = \{f \in W_1^{2,0}(G) : f^*|_E = 0 \text{ q.e.}\}.$

**5.2. Traces.** A result of [16], p. 182, shows that the trace operator

$$T : W_1^{2,0}(G) \rightarrow B_2, \quad Tf = f^*|_{\mathbb{T}}$$

is a well defined, continuous, surjective operator with, by Proposition 5.1,

$$\ker(T) = W_1^{2,0}(G \setminus \mathbb{T}).$$



**Remark.** By [18], one can also define the (continuous) trace operator

$$\mathrm{tr} : W_1^{2,0}(G) \rightarrow B_2$$

$$(\mathrm{tr} f)(\zeta) = \lim_{r \uparrow 1} f(r\zeta)$$

and notice that this limit exists a.e.  $|d\zeta|$  and in  $L^2(\mathbb{T}, |d\zeta|)$ . Also notice that  $T(\phi) = \mathrm{tr}(\phi)$  for all  $\phi \in C_0^\infty(G)$ . Thus  $Tf = \mathrm{tr} f$  a.e.  $|d\zeta|$  for all  $f \in W_1^{2,0}(G)$ .

Thus (as before)  $T$  will induce

$$\tilde{T} : W_1^{2,0}(G)/W_1^{2,0}(G \setminus \mathbb{T}) \rightarrow B_2, \quad \tilde{T}[h] = h^*|_{\mathbb{T}}$$

and  $\tilde{T}$  will be continuous and invertible. So (as before) we define

$$J : L_a^2(G)^\perp / L_a^2(G \setminus \mathbb{T})^\perp \rightarrow B_2$$

by  $J = \tilde{T} \circ \tilde{C}$  to obtain

$$J[g](\zeta) = -\frac{1}{\pi} \int_G \frac{g(z)}{z - \zeta} dA(z) \quad \text{q.e.}$$

and  $JR = M_\zeta J$  (see [2] for details). Thus there is a one-to-one correspondence between the invariant subspaces  $\chi_G \in \mathcal{M} \subset L_a^2(G \setminus \mathbb{T})$  and the invariant subspaces of  $B_2$ . One also shows (as before) that for an invariant subspace  $\chi_{\mathbb{D}}, \chi_G \in \mathcal{M} \subset L_a^2(G \setminus \mathbb{T})$  and  $g \in \mathcal{M}^\perp$ ,

$$J[g](\zeta) = -\frac{1}{\pi} \int_{G \setminus \mathbb{D}} \frac{g(z)}{z - \zeta} dA(z)$$

and so  $J[g] \in AB_2$ . So, as before, there is a one-to-one correspondence between the invariant subspaces  $\chi_{\mathbb{D}}, \chi_G \in \mathcal{M}$  and the invariant subspaces of  $AB_2$ .

**5.3. Zero Sets.** For a quasi-closed set  $E \subset \mathbb{T}$ , we can find a sequence of closed sets  $F_1 \subset F_2 \subset \cdots \subset E$  with  $C_2(F_n) \rightarrow C_2(E)$ . Since  $L_a^2(G \setminus F_n)$  increases with  $n$ , we can define the invariant subspace

$$(5.2) \quad \mathcal{M}(E) = \overline{\bigcup_n L_a^2(G \setminus F_n)}^{L^2}.$$

One can prove the following basic facts about  $\mathcal{M}(E)$  [23]:

**Proposition 5.2.** For quasi-closed sets  $E, F \subset K$

- (1)  $\mathcal{M}(E)$  is independent of the choice of  $\{F_n\}$ .
- (2)  $\mathcal{M}(E) \subset \mathcal{M}(F) \Leftrightarrow C_2(E \setminus F) = 0$ .
- (3)  $\mathcal{M}(E) = \mathcal{M}(F) \Leftrightarrow C_2(E \Delta F) = 0$ .

**Remark.** There are quasi-closed sets  $E \subset \mathbb{T}$  for which  $\mathcal{M}(E)$  cannot be written as  $L_a^2(G \setminus F)$  for any closed  $F \subset K$  [23], Proposition 4.3.

One also notes that

$$(5.3) \quad W_2(E) \equiv C(\mathcal{M}(E)^\perp) = \bigcap_n W_1^{2,0}(G \setminus F_n)$$

and by Proposition 2.1,  $f \in W_1^{2,0}(G)$  belongs to  $W_2(E)$  if and only if  $f^* = 0$  quasi-everywhere on  $E$ . From this one has

$$(5.4) \quad \begin{aligned} J(\mathcal{M}(E)^\perp / L_a^2(G \setminus \mathbb{T})^\perp) &= T(C\mathcal{M}(E)^\perp) = T(W_2(E)) \\ &= \{f \in B_2 : f^*|_E = 0 \text{ q.e.}\}. \end{aligned}$$

**Remark.** The subspace  $B_{2,E}(\mathbb{T}) \equiv \{f \in B_2 : f^*|_E = 0 \text{ q.e.}\}$  can be described in several equivalent ways. If  $f \in B_2$ , then (as mentioned in the introduction, see also [20]) the radial limit of its harmonic extension exists q.e. and is a quasi-continuous function on  $\mathbb{T}$ . Since  $f^*$  is also a quasi-continuous function on  $\mathbb{T}$  which equals the radial limit function a.e.  $|d\zeta|$ , then [20], Proposition 2.1 (c),

$$f^*(\zeta) = \lim_{r \rightarrow 1} f(r\zeta) \quad \text{q.e.}$$

Thus we have that  $Z(f)$  is quasi-closed (an easy exercise using the definition of quasi-continuity) and

$$B_{2,E}(\mathbb{T}) = \{f \in B_2 : C_2(E \setminus Z(f)) = 0\}$$

from which  $AB_{2,E} = B_{2,E} \cap H^2$ .

**6. Invariant subspaces for  $p = 2$ .** For an inner function  $I$  and a quasi-closed set  $E \subset \mathbb{T}$ , recall that

$$\mathcal{F}_{E,I} = IH^2 \cap AB_{2,E}$$

is a (closed) invariant subspace of  $AB_2$ . We now identify  $\mathcal{M}_{\mathcal{F}_{E,I}}$ .

**Theorem 6.1.**

$$\mathcal{M}_{\mathcal{F}_{E,I}} = \mathcal{M}(E) \bigvee \left\{ \frac{\chi_{G \setminus \mathbb{D}}}{\phi} L_a^2(G) : \phi \in V(I) \right\}.$$

*Proof.* The proof is nearly identical to the proof of Theorem 4.1, except for some technicalities. We first show that

$$(6.1) \quad \mathcal{M}(E) = \overline{\bigcup_n L_a^2(G \setminus F_n)}^{L^2} \subset \mathcal{M}_{\mathcal{F}_{E,I}}.$$

Fix  $n$  and let  $h \in L_a^2(G \setminus F_n)$  and  $f \in W_1^{2,0}(G)$  with  $Tf \in IH^2 \cap B_{2,E}$ . Then by Proposition 2.1,  $f \in W_1^{2,0}(G \setminus F_n)$  and so by Havin's Lemma, Lemma 2.2,

$$\langle \bar{\partial} f, h \rangle = 0.$$

Thus by the definition of  $\mathcal{M}(E)$ , we have (6.1).

By (3.3), notice that

$$\mathcal{M}_{\mathcal{F}_{E,I}} = \left( \bar{\partial} \{ f \in W_1^{2,0}(G) : Tf \in IH^2 \cap AB_{2,E} \} \right)^\perp.$$

So let  $f \in W_1^{2,0}(G)$  with  $Tf \in IH^2 \cap AB_{2,E}$ . Then for all  $\phi \in V(I)$ ,

$$\int_G \bar{\partial} f \frac{\chi_{G \setminus \mathbb{D}}}{\phi} dA = \int_{G \setminus \mathbb{D}} \frac{\bar{\partial} f}{\phi} dA = \lim_{\varepsilon \rightarrow 0} \int_{G \setminus \{|z| < 1+\varepsilon\}} \bar{\partial} \left( \frac{f}{\phi} \right) dA,$$

which by Green's theorem [31], p. 54, equals

$$\lim_{\varepsilon \rightarrow 0} -\frac{1}{2i} \int_{|z|=1+\varepsilon} \frac{f}{\phi} dz.$$

By [18],  $f((1+\varepsilon)\zeta) \rightarrow f(\zeta)$  in  $L^2(\mathbb{T}, |d\zeta|)$ . Thus using the fact that  $Tf/\phi \in H^2$ , the above becomes

$$-\frac{1}{2i} \int_{\mathbb{T}} \frac{Tf}{\phi} dz = 0,$$

hence  $\chi_{G \setminus \mathbb{D}}/\phi \in \mathcal{M}_{\mathcal{F}_{E,I}}$ . So, by the invariance of  $\mathcal{M}_{\mathcal{F}_{E,I}}$ , the density of polynomials in  $L_a^2(G)$ , and (6.1),

$$\mathcal{M}(E) \bigvee \left\{ \frac{\chi_{G \setminus \mathbb{D}}}{\phi} L_a^2(G) : \phi \in V(I) \right\} \subset \mathcal{M}_{\mathcal{F}_{E,I}}.$$

By taking annihilators and then  $C = \bar{\partial}^{-1}$  we get (using (5.3))

$$C\mathcal{M}_{\mathcal{F}_{E,I}}^\perp \subset W_2(E) \cap C \left( \bigcap \left\{ \left( \frac{\chi_{G \setminus \mathbb{D}}}{\phi} L_a^2(G) \right)^\perp : \phi \in V(I) \right\} \right).$$

Let  $g = C(\bar{\partial}g)$  belong to the right hand side of the above. Then  $g^*|_E = 0$  q.e. and

$$\int_{G \setminus \mathbb{D}} \bar{\partial}g \frac{z^n}{I} dA = 0 \quad \forall n \in \mathbb{N} \cup \{0\}.$$

As before (using Green's theorem)

$$\int_{\mathbb{T}} \frac{g}{I} z^n dz = 0 \quad \forall n \in \mathbb{N} \cup \{0\}$$

and so  $(Tg)/I \in H^2$  which means  $Tg \in IH^2 \cap AB_{2,E}$ . Thus  $g \in C\mathcal{M}_{\mathcal{F}_{E,I}}^\perp$  and we are done.  $\square$

Using a similar proof as in Corollary 4.2, one can prove the following corollary:

**Corollary 6.2.** *If the inner function  $I$  is the least common multiple of the inner functions  $I_1$  and  $I_2$ , then*

$$\mathcal{M}_{\mathcal{F}_{E,I}} = \mathcal{M}(E) \bigvee \frac{\chi_{G \setminus \mathbb{D}}}{I_1} L_a^2(G) \bigvee \frac{\chi_{G \setminus \mathbb{D}}}{I_2} L_a^2(G).$$

As in the  $1 < p < 2$  case we have  $\mathcal{M}_{\mathcal{F}_{E,I}} = L_a^2(G \setminus \mathbb{T})$  if and only if  $\mathcal{F}_{E,I} = 0$ . However, understanding when  $\mathcal{F}_{E,I}$  is non-trivial is more complicated and is yet unknown. In fact, understanding the zero sets for the Dirichlet space (i.e. when  $\mathcal{F}_{E,1} \neq 0$ ) remains an open problem [10]. For example, there are Blaschke products which divide Dirichlet functions and whose zeros accumulate near every point of  $\mathbb{T}$ , in stark contrast to the  $AB_q$  ( $q > 2$ ) case where the zeros must accumulate on a Carleson set.

By [22], it is known that every invariant subspace  $\mathcal{F}$  of  $AB_2$  is of the form

$$\mathcal{F} = IH^2 \cap [f],$$

where  $I$  is an inner function,  $f \in AB_2$  is outer and  $[f] = \text{span}\{\zeta^n f : n = 0, 1, 2, \dots\}$ . Certainly  $\mathcal{F} \subset \mathcal{F}_{Z(f),I}$  and so by (3.3)

$$\mathcal{M}_{\mathcal{F}} \supset \mathcal{M}_{\mathcal{F}_{Z(f),I}}.$$

It is a conjecture that  $[f] = AB_{2,Z(f)}$  and hence we will have equality above, thus giving us a complete characterization of the invariant subspaces  $\chi_{\mathbb{D}}, \chi_G \in \mathcal{M} \subset L^2_a(G \setminus \mathbb{T})$ . For certain outer functions  $f$ ,  $[f] = AB_{2,Z(f)}$ , but the general question still remains open.

**7.  $p > 2$ .** We mention that the results in the previous section have generalizations to  $p > 2$ . The techniques are exactly the same except for some technicalities which we mention now.

For  $p \geq 2$  the appropriate space to look at is  $B_q$  and the capacity used is the  $C_q$  capacity (defined in an analogous way). The capacity theory is the same and the trace operator  $T$  is defined as before.

If  $1 < q \leq 2$  and  $f \in AB_q = D_q$ , it is known [8] that the radial limit

$$\lim_{r \rightarrow 1} f(r\zeta)$$

exist everywhere except possibly on a set of  $q$ -Bessel capacity (equivalently the  $C_q^*$  capacity) zero. As is the  $q = 2$  case, we define the set

$$Z(f) = \{\zeta \in \mathbb{T} : \lim_{r \rightarrow 1} f(r\zeta) = 0\}.$$

For a quasi-closed set (with respect to the  $C_q$  capacity)  $E \subset \mathbb{T}$  we define

$$AB_{q,E} = \{f \in AB_q : C_q(E \setminus Z(f)) = 0\}.$$

A result of Carleson [7] says that  $AB_{2,E}$  is a closed subspace of  $AB_2$ . This next result (which is known but we could not find a proof) says the same for  $AB_{q,E}$ .

**Proposition 7.1.** *For  $1 < q \leq 2$  and a set  $E \subset \mathbb{T}$ ,  $AB_{q,E}$  is a closed subspace of  $AB_q$ .*

*Proof.* It is known (by observing that  $W_1^q$  is the same as the space of Bessel potentials  $L_1^q$ ) that if  $\{f_i\}$  is a Cauchy sequence of quasi-continuous functions in  $AB_q = D_q$  (i.e. quasi-everywhere defined on  $\mathbb{D}$  and quasi-continuous) then there is a quasi-continuous  $f \in AB_q$  and a subsequence  $f_{i_j} \rightarrow f$  quasi-everywhere (see [20], Proposition 2.1 for a proof in a slightly different setting).

Thus for  $f \in AB_q$  we define  $\tilde{f}$  q.e. on  $\mathbb{D}$  by setting  $\tilde{f}$  to be  $f(z)$  for  $z \in \mathbb{D}$  and  $\tilde{f}(\zeta)$  to be the radial limit of  $f$  at  $\zeta$  for  $\zeta \in \mathbb{T}$ . For  $0 < r < 1$  define  $f_r(z) = f(rz)$  and notice that  $f_r$  is continuous on  $\mathbb{D}$  and  $f_r \rightarrow f$  in  $L^q$ -Dirichlet norm. Thus by the above fact,  $\tilde{f}$  is quasi-everywhere equal to a quasi-continuous function, making  $\tilde{f}$  quasi-continuous on  $\mathbb{D}$ .

So if  $\{f_n\}$  is a Cauchy sequence in  $AB_{q,E}$  then  $\{\tilde{f}_n\}$  is a Cauchy sequence of quasi-continuous functions in  $AB_{q,E}$  and by the above fact, the limit function must vanish q.e. on  $E$ . Thus  $AB_{q,E}$  is closed.  $\square$

For  $f \in AB_q$  we have that  $f^*$  and  $\tilde{f}$  are quasi-continuous functions on  $\bar{\mathbb{D}}$  with  $f^* = \tilde{f}$  a.e. By [3], Theorem 2 (iii),  $f^* = \tilde{f}$  q.e. As before, one defines  $\mathcal{M}(E)$  and one has (using the above and (5.4)) that

$$J(\mathcal{M}(E)^\perp / L_a^p(G \setminus \mathbb{T})^\perp) = AB_{q,E}.$$

For an inner function  $I$  and a quasi-closed set  $E \subset \mathbb{T}$  define

$$\mathcal{F}_{E,I} = IH^q \cap AB_{q,E}$$

and notice that this is a closed subspace of  $AB_q$ . Using the same proof as in the  $q = 2$  case and the above, one proves that

$$\mathcal{M}_{\mathcal{F}_{E,I}} = \mathcal{M}(E) \bigvee \left\{ \frac{\chi_{G \setminus \mathbb{D}}}{\phi} L_a^p(G) : \phi \in V(I) \right\}.$$

One also proves, in exactly the same way as before, that if the inner function  $I$  is the least common multiple of the inner functions  $I_1$  and  $I_2$ , then

$$\mathcal{M}_{\mathcal{F}_{E,I}} = \mathcal{M}(E) \bigvee \frac{\chi_{G \setminus \mathbb{D}}}{I_1} L_a^p(G) \bigvee \frac{\chi_{G \setminus \mathbb{D}}}{I_2} L_a^p(G).$$

**8. Codimension.** If  $1 < p \leq 2$  and  $S : L_a^p(G \setminus \mathbb{T}) \rightarrow L_a^p(G \setminus \mathbb{T})$  is  $(Sf)(z) = zf(z)$ , then for an invariant subspace  $\mathcal{M}$  and  $\lambda \in G \setminus \mathbb{T}$ ,  $(S - \lambda)|_{\mathcal{M}}$  is a semi-Fredholm operator and

$$-\text{index}((S - \lambda)|_{\mathcal{M}}) = \dim(\mathcal{M}/(z - \lambda)\mathcal{M})$$

is constant on the components of  $G \setminus \mathbb{T}$  [19], Lemma 2.1, and is called the *codimension* on the component of  $G \setminus \mathbb{T}$ . In [2] they prove the following formula:

$$(8.1) \quad \dim(\mathcal{M}/\mathcal{M}(z - \lambda)) = 1 + \dim(\mathcal{F}_{\mathcal{M}}/(\zeta - \lambda)\mathcal{F}_{\mathcal{M}}),$$

where  $\mathcal{F}_{\mathcal{M}}$  is the unique invariant subspace of  $B_q$  that corresponds to  $\mathcal{M}$  via the operator  $J$ .

**Theorem 8.1.** *If  $1 < p \leq 2$  and  $\chi_G, \chi_{\mathbb{D}} \in \mathcal{M} \subset L_a^p(G \setminus \mathbb{T})$  is a non-trivial invariant subspace, then for  $\lambda \in G \setminus \mathbb{T}$ ,*

$$\dim(\mathcal{M}/(z - \lambda)\mathcal{M}) = \begin{cases} 1 & \text{if } |\lambda| > 1 \\ 2 & \text{if } |\lambda| < 1 \end{cases}$$

*Proof.* If  $\lambda \in G \setminus \mathbb{T}$  with  $|\lambda| > 1$ , then  $M_{\zeta-\lambda}$  is an invertible operator on  $\mathcal{F}_{\mathcal{M}}$  and so  $(\zeta - \lambda)\mathcal{F}_{\mathcal{M}} = \mathcal{F}_{\mathcal{M}}$ . From (8.1) we get  $\dim(\mathcal{M}/(z - \lambda)\mathcal{M}) = 1$ .

Since  $\mathcal{F}_{\mathcal{M}}$  is a non-trivial invariant subspace of  $AB_q$  then by a result of [19] ( $q > 2$ ) and [21] ( $q = 2$ ),  $\dim(\mathcal{F}_{\mathcal{M}}/\zeta\mathcal{F}_{\mathcal{M}}) = 1$ . Thus by (8.1),  $\dim(\mathcal{M}/z\mathcal{M}) = 2$ . Since the codimension is constant on the components of  $G \setminus \mathbb{T}$  we are done.  $\square$

We mention that in general the invariant subspaces  $\mathcal{F}$  of  $B_2$  can be quite complicated, thus making the invariant subspaces  $\chi_G \in \mathcal{M} \subset L_a^2(G \setminus \mathbb{T})$  difficult to describe. The invariant subspaces  $\mathcal{F}$  with  $\zeta\mathcal{F} = \mathcal{F}$  have been completely characterized [20] as  $\mathcal{F} = B_{2,E}(\mathbb{T})$  for some quasi-closed  $E \subset \mathbb{T}$  and thus by (5.4)  $\mathcal{M}_{\mathcal{F}} = \mathcal{M}(E)$ . The invariant subspaces  $\mathcal{F}$  with  $\zeta\mathcal{F} \neq \mathcal{F}$  (such subspaces are called *simply invariant*) are quite complicated. In fact, recall from the introduction that the invariant subspace  $\mathcal{M}_n$  of the Bergman space, (1.2), has  $\dim(\mathcal{M}_n/z\mathcal{M}_n) = n$ . Thus by (8.1)

$$\dim(\mathcal{F}_{\mathcal{M}_n}/\zeta\mathcal{F}_{\mathcal{M}_n}) = n,$$

(see [24] for a specific example) which is in stark contrast to the analytic Dirichlet space  $AB_2$  where the codimension of any non-zero invariant subspace is always one.

## REFERENCES

- [1] R. A. ADAMS, *Sobolev Spaces*, Academic Press, New York, 1975.
- [2] A. ALEMAN, S. RICHTER AND W. T. ROSS, *Bergman spaces on disconnected domains*, (Preprint).
- [3] T. BAGBY, *Quasi topologies and rational approximation*, J. Funct. Anal. **10** (1972), 259-268.
- [4] H. BERCOVICI, C. FOIAŞ, AND C. PEARCY, *Dual algebras with applications to invariant subspaces and dilation theory*, CBMS Regional Conf. Ser. in Math., no. 56, Amer. Math. Soc., Providence, RI., 1985.
- [5] O. V. BESOV, *On embedding and extension theorems for some function classes*, Trudy Mat. Inst. Steklov **60** (1960), 42-81 (Russian). Translation: Amer. Math. Soc. Transl. **40** (1964), 85-126.
- [6] A. BEURLING, *Ensembles exceptionnels*, Acta Math. **72** (1940), 1-13.
- [7] L. CARLESON, *Sets of uniqueness for functions regular in the unit circle*, Acta Math. **87** (1952), 325-345.
- [8] L. CARLESON, *Selected problems on exceptional sets*, D. Van Nostrand, Princeton, New Jersey, 1967.
- [9] L. CARLESON, *A representation formula for the Dirichlet integral*, Math. Z. **73** (1960), 190-196.
- [10] L. CARLESON, *On the zeros of functions with bounded Dirichlet integrals*, Math. Z. **56** (1952), 289-295.
- [11] H. FEDERER AND W. ZIEMER, *The Lebesgue set of a function whose distribution derivatives are  $p$ -th power summable*, Indiana Univ. Math. J. **22** (1972), 139-158.
- [12] V. P. HAVIN, *Approximation in the mean by analytic functions*, Dokl. Akad. Nauk SSSR **178** (1968), 1025-1028 (Russian). Translation: Soviet Math. Dokl. **9** (1968), 245-248.
- [13] V. P. HAVIN, *The factorization of analytic functions that are smooth up to the boundary*, Zap. Nauch. Sem. Leningrad. Otdel. Math. Inst. Steklov. (LOMI) **22** (1971), 202-205.

- [14] L. I. HEDBERG, *Non-linear potentials and approximation in the mean by analytic functions*, Math. Z. **129** (1972), 299–319.
- [15] K. HOFFMAN, *Banach Spaces of Analytic Functions*. Dover Publications, New York, 1988.
- [16] A. JONSSON AND H. WALLIN, *Function spaces on subsets of  $\mathbb{R}^n$* , Mathematical Reports, Vol. 2, Harwood Academic Publishers, London-Paris-Utrecht-New York, 1984.
- [17] S. KHRUSHCHEV AND V. PELLER, *Hankel operators, best approximation, and stationary Gaussian processes*, Russian Math. Surveys **37** (1982), 61–144.
- [18] P. I. LIZORKIN, *Boundary properties of functions from ‘weighted’ classes*, Dokl. Akad. Nauk SSSR **132** (1960), 514–517 (Russian). Translation: Soviet Math. Dokl. **1** (1960), 589–593.
- [19] S. RICHTER, *Invariant subspaces in Banach spaces of analytic functions*, Trans. Amer. Math. Soc. **304** (1987), 585–616.
- [20] S. RICHTER, W. T. ROSS, AND C. SUNDBERG, *Hyperinvariant subspaces of the harmonic Dirichlet space*, J. Reine Angew. Math. **448** (1994), 1–26.
- [21] S. RICHTER AND A. SHIELDS, *Bounded analytic functions in the Dirichlet space*, Math. Z. **198** (1988), 151–159.
- [22] S. RICHTER AND C. SUNDBERG, *Multipliers and invariant subspaces in the Dirichlet space*, Jour. Op. Theory (to appear).
- [23] W. T. ROSS, *Invariant subspaces of Bergman spaces on slit domains*, Bull. London Math. Soc. (to appear).
- [24] W. T. ROSS, *Invariant subspaces of the harmonic Dirichlet space with large codimension*, Proc. Amer. Math. Soc. (to appear).
- [25] W. RUDIN, *The closed ideals in an algebra of analytic functions*, Can. J. Math. **9** (1957), 426–434.
- [26] N. A. SHIROKOV, *Closed ideals of algebras of type  $B_{p,q}^\alpha$* , Izv. Akad. Nauk SSSR, Math. **46** (1982), 1316–1333 (Russian). Translation: Math. USSR Izvestiya **21** (1983), 585–600.
- [27] N. A. SHIROKOV, *Analytic Functions Smooth up to the Boundary*, Lecture Notes in Math. **1312**, Springer-Verlag, New York/Berlin, 1988.
- [28] E. M. STEIN, *Singular Integrals and Differentiability Properties of Functions*. Princeton Univ. Press, Princeton, New Jersey, 1970.
- [29] B. A. TAYLOR AND D. L. WILLIAMS, *Zeros of Lipschitz functions in the unit disc*, Mich. Math. J. **18** (1971), 129–139.
- [30] B. A. TAYLOR AND D. L. WILLIAMS, *Ideals in rings of analytic functions with smooth boundary values*, Can. J. Math. **22** (1970), 1266–1283.
- [31] I. N. VEKUA, *Generalized Analytic Functions*, Addison-Wesley, Reading, Mass., 1962.

This research was supported in part from a grant from the National Science Foundation.

Department of Mathematics  
University of Richmond  
Richmond, Virginia 23173

E-mail: rossb@@mathcs.urich.edu

Received: April 7th, 1994.